arXiv:1103.0107v1 [math.CA] 1 Mar 2011

Mixed means of commutators of central integral means and CMO

Shunchao Long and Jian Wang

In this paper we obtain some mixed means and weighted L^p estimates for the commutators generating r order central integral means operators with CMO functions.

1.INTRODUCTION

Let T be a sublinear operator and $b(x)$ a function, the commutator $[T,b]$ is defined by

$$[T,b]f(x)=T(fb)(x)-b(x)Tf(x).$$

A famous result of Coifman, Rochberg and Weiss stated that the commutators of Calderon-Zygmund singular integral operators and BMO functions are bounded on L^p for $1<p<\infty$ (see [5]). Since then, the L^p estimates and applications of these type commutators were studied by many authors (see [2,21,19,4] and [3,15,20,23]).

Recently the authors of this paper proved the L^p-boundedness of the commutators of Hardy operators and one-side CMO functions for $1<p<\infty$ in [18]. And this result was extended in [16,11]. In this paper we establish some mixed means estimates and weighted L^p-estimates of the commutators of central integral means and CMO functions, and this extends the boundedness results in [18,11].

Let $1\le p<\infty$. Denote

$$\sup_{B\subset\mathbf{R}^n}\left(\frac{1}{|B|}\int_B|b(x)-b_B|^pdx\right)^{1/p}=\begin{cases}\|b\|_{BMO^p}, & \text{if } B \text{ are arbitrary balls,}\\ \|b\|_{CMO^p}, & \text{if } B \text{ are balls centraled at origin,}\end{cases}$$

where $b_B=\frac{1}{|B|}\int_B b(x)dx$. If $\|b\|_{BMO^p}<\infty$, we say $b\in BMO^p$, the Bounded Mean Oscillation. It is well-known that $BMO^p=BMO^1=BMO$ for all $1\le p<\infty$. If $\|b\|_{CMO^p}<\infty$, we say $b\in CMO^p$, the Central Mean Oscillation. Obviously, if $1\le p<q<\infty$, then $CMO^q\subset CMO^p$, and $\|b\|_{CMO^p}\le\|b\|_{CMO^q}$.

We say $b\in CMO$, if $\|b\|_{CMO}=\sup_{1\le p<\infty}\|b\|_{CMO^p}<\infty$.

The spaces CMO^p bear a simple relationship with BMO: $g\in BMO$ precisely when g and all of its translates belong to CMO^p uniformly a.e., so

$$\text{the classical } BMO \text{ space } \subset CMO^p \text{ for all } 1\le p<\infty.$$

Many precise analogies exist between CMO^p and BMO from the point of view of real Hardy spaces, for example, the duality: $CMO^{p'}, p' = p/(p-1)$, is the dual of the Beurling-Herz-Hardy spaces $HA^p, 1 \leq p < \infty$, that is then analogous to the $H^1 \leftrightarrow BMO$, (see [11-12]); the constructive decomposition (see [17]); and so on.

If $R > 0$, denote $B(R) = B(0, R)$ be the ball in $\mathbf{R}^n$ centered at origin and of radius R. Let $r, \alpha \in \mathbf{R}, f \in L^r_{loc}(\mathbf{R}^n, |x|^{n(\alpha-1)})$, the central integral mean of order r, with the power weight, of f, be defined by

$$M_r(f,\alpha)(|y|) = \left[\frac{1}{|B(|y|)|^\alpha}\int_{B(|y|)} |B(|x|)|^{\alpha-1}|f(x)|^r dx\right]^{1/r},$$

and the companion mean of order r, with the power weight, of f, by

$$M^*_r(f,\alpha)(|y|) = \left[\frac{1}{|B(|y|)|^\alpha}\int_{\mathbf{R}^n\setminus B(|y|)} |B(|x|)|^{\alpha-1}|f(x)|^r dx\right]^{1/r}.$$

The properties and applications of these types of integral means can be found in many literatures. Firstly, the limits of $(M_2(f,1)(y))^2 = (1/2y)\int_{-y}^{y}|f|^2 (y > 0$, the one-dimensional case), were used to study the almost periodic functions, and the spectrum and ergodicity of sample paths of certain stochastic processes by Wiener in [22]. Secondly, the functions spaces of bounded integral mean of r order, introduced firstly by Beurling,

$$B^r = B^{r,\infty} = \{f : M_r(f,1)(|y|) \in L^\infty\},$$

(both homogeneous spaces ($|y| > 0$) and non-homogeneous spaces ($|y| > 1$)), together with their corresponding Beurling algebra A^r and the Hardy space HA^r [cf, 1, 12, 6] had rich contents; Thirdly, the Hardy type inequalities associating with $M_1(f,1)(|y|)$ and $M^*_1(f,1)(|y|)$ generalized the classical ones to n-dimension ball [9-10, 7]. And the mixed means inequalities were used to derive the generalizing Hardy and Levin-Cochran-Lee type inequalities in [7] (see also [8]).

We state the mixed means inequalities as following:

Theorem 1 ([7: Theorem 5]). Let $r, s, R, \alpha, \gamma \in \mathbf{R}$, and let $r < s, r, s \neq 0, R > 0 (f \neq 0$ in the case of $r < 0)$. Then,

$$M_s((M_r(f,\alpha),\gamma)(R) \leq M_r((M_s(f,\gamma),\alpha)(R), \tag{1}$$

$$M^*_s((M^*_r(f,\alpha),\gamma)(R) \leq M^*_r((M^*_s(f,\gamma),\alpha)(R). \tag{2}$$

From Theorem 1, we can obtain the L^p-boundedness estimates of these types of integral means above as following.

Theorem 2. Let $r, s, \alpha, \gamma \in \mathbf{R}$, and let $r < s, r \neq 0, s > 0 (f \neq 0$ in the case of $r < 0)$. Then,

$$\int_{\mathbf{R}^n} |B(|y|)|^{\gamma-1}(M_r(f,\alpha)(|y|))^s dy \leq \frac{1}{(\alpha-\gamma r/s)^{s/r}}\int_{\mathbf{R}^n} |B(|y|)|^{\gamma-1}|f(y)|^s dy \tag{3}$$

if $\alpha - \gamma r/s > 0$, and

$$\int_{\mathbf{R}^n} |B(|y|)|^{\gamma-1}(M_r^*(f,\alpha)(|y|))^s dy \le \frac{1}{(\gamma r/s-\alpha)^{s/r}} \int_{\mathbf{R}^n} |B(|y|)|^{\gamma-1}|f(y)|^s dy \qquad (4)$$

if $\alpha - \gamma r/s < 0$.

Proof . For $y \in B(R)$, $\int_{B(|y|)} |B(|x|)|^{\gamma-1}|f(x)|^s dx \le \int_{B(R)} |B(|x|)|^{\gamma-1}|f(x)|^s dx$ and $\int_{B(R)} |B(|y|)|^{\alpha-1-\gamma r/s} dy = \frac{1}{\alpha-\gamma r/s}|B(R)|^{\alpha-\gamma r/s}$ when $\alpha - \gamma r/s > 0$, and using (1), we have

$$\begin{aligned}
&\int_{B(R)} |B(|y|)|^{\gamma-1}(M_r(f,\alpha)(|y|))^s dy \\
&\le |B(R)|^{\gamma-s\alpha/r} \left[\int_{B(R)} |B(|y|)|^{\alpha-1} \left(\frac{1}{|B(|y|)|^\gamma} \int_{B(|y|)} |B(|x|)|^{\gamma-1}|f(x)|^s dx\right)^{r/s} dy\right]^{s/r} \\
&\le \frac{1}{(\alpha-\gamma r/s)^{s/r}} \int_{B(R)} |B(|x|)|^{\gamma-1}|f(x)|^s dx,
\end{aligned}$$

(3) follows by taking the $\lim_{R\to\infty}$. And (4) is the consequence of (2), derived by the same technique as (3) from (1), except for taking the $\lim_{R\to 0}$.

Remark It is easy to see that $L^\infty \subset B^{r,\infty}$ for $0 < r < \infty$ (see also [6,12-13]). Let

$$B^{r,s}(\alpha,\gamma) = \{f : M_r(f,\alpha)(|y|) \in L^s_{|x|^{n\gamma}}\}.$$

Then r, s, α, γ are under the conditions of Theorem 2 and $\alpha - \gamma r/s > 0$. Thus, we have

$$L^s_{|x|^{n\gamma}} \subset B^{r,s}(\alpha,\gamma).$$

Further, we can obtain the boundedness estimates of the commutators generating r order central integral means operators and CMO functions.

From the point of view on Hardy spaces, the central integral means bear some relationships with CMO^p. In fact, $(A^p)^* = B^{p'}$ and $(HA^p)^* = CMO^{p'}$.

2. COMMUTATOR THEOREMS

Let $r > 0, \alpha \in \mathbf{R}, f \in L^r_{loc}(\mathbf{R}^n, |x|^{n(\alpha-1)})$, and b be local integral functions on $\mathbf{R}^n$. We define the integral mean commutators of order r, with the power weight, by

$$M_{r,b}(f,\alpha)(|y|) = \left[\frac{1}{|B(|y|)|^\alpha} \int_{B(|y|)} |B(|x|)|^{\alpha-1}(|b(x)-b(y)||f(x)|)^r dx\right]^{1/r},$$

and the companion mean commutators of order r, with the power weight, by

$$M^*_{r,b}(f,\alpha)(|y|) = \left[\frac{1}{|B(|y|)|^\alpha} \int_{\mathbf{R}^n\setminus B(|y|)} |B(|x|)|^{\alpha-1}(|b(x)-b(y)||f(x)|)^r dx\right]^{1/r}.$$

When $r \ge 1$, by the Minkowski inequality, it is easy to calculate that,

$$|[M_r(\bullet,\alpha), b]f(|y|)| \stackrel{def}{=} |M_r(bf,\alpha)(|y|) - b(|y|)M_r(f,\alpha)(|y|)| \le M_{r,b}(f,\alpha)(|y|),$$

$|[M_r^*(\bullet,\alpha),b]f(|y|)| \overset{def}{=} |M_r^*(bf,\alpha)(|y|) - b(|y|)M_r^*(f,\alpha)(|y|)| \le M_{r,b}^*(f,\alpha)(|y|)$.

Mixed-means inequalities of commutators of integral means and CMO functions:

Theorem 3. Let $r,s,R,\alpha,\gamma \in \mathbf{R}, s > r > 0, R > 0$. Suppose that $b \in CMO$ and f is local bounded functions. Then,

$$M_s((M_{r,b}(f,\alpha),\gamma)(R) \le c_1\|b\|_{CMO}M_r((M_s(f,\gamma),1)(R) \tag{5}$$

if $\alpha > 1$; and

$$M_s^*((M_{r,b}^*(f,\alpha),\gamma)(R) \le c_1\|b\|_{CMO}M_r^*((M_s^*(f,\gamma),1)(R) \tag{6}$$

if $\alpha < 1$, where $c_1 = [2^{n|\alpha|}2^{n|\alpha-1|}3^r \times 4 \times 2^{2nr}\sum\limits_{k=0}^{\infty} 2^{-kn|\alpha-1|}k^r]^{1/r}$.

Weighted L^p-estimates of commutators of integral means and CMO functions:

Theorem 4. Let $r,s,\alpha,\gamma \in \mathbf{R}, s > r > 0; b \in CMO, f$ is local bounded functions, then,

$$\int_{\mathbf{R}^n} |B(|y|)|^{\gamma-1}(M_{r,b}(f,\alpha)(|y|))^s dy \le c_2\|b\|_{CMO}^s \int_{\mathbf{R}^n} |B(|y|)|^{\gamma-1}|f(y)|^s dy, \tag{7}$$

if $\alpha > 1$ and $\gamma < s/r$, and

$$\int_{\mathbf{R}^n} |B(|y|)|^{\gamma-1}(M_{r,b}^*(f,\alpha)(|y|))^s dy \le c_2\|b\|_{CMO}^s \int_{\mathbf{R}^n} |B(|y|)|^{\gamma-1}|f(y)|^s dy, \tag{8}$$

if $\alpha < 1$ and $\gamma > s/r$, where $c_2 = \frac{1}{|1-\gamma r/s|^{s/r}}[2^{n|\alpha|}2^{n|\alpha-1|}3^r \times 4 \times 2^{2nr}\sum\limits_{k=0}^{\infty} 2^{-kn|\alpha-1|}k^r]^{s/r}$.

Proof of Theorem 3 Let us prove (5). Let $2^{N-1} < R < 2^N$, denote

$$B_i = \begin{cases} B(2^i), & \text{if } i \le N-1, \\ B(R), & \text{if } i = N, \end{cases}, C_i = B_i \setminus B_{i-1}, i = -\infty, ..., N; \tag{9}$$

if $x \in C_i$, denote

$$\overline{B_j} = \begin{cases} B_j, & \text{if } j \le i-1, \\ B(|x|), & \text{if } j = i, \end{cases}, \overline{C_j} = \overline{B_j} \setminus \overline{B_{j-1}}, j = -\infty, ..., i. \tag{10}$$

Then, we have

$$h(x) = [(M_{r,b}(f,\alpha)(|x|)]^r = \frac{1}{|B(|x|)|^\alpha}\sum_{j=-\infty}^{i}\int_{\overline{C_j}} |B(|y|)|^{\alpha-1}(|b(x)-b(y)||f(y)|)^r dy,$$

and

$$[M_s((M_{r,b}(f,\alpha),\gamma)(R)]^s = \frac{1}{|B(R)|^\gamma}\sum_{i=-\infty}^{N}\int_{C_i} |B(|x|)|^{\gamma-1}\,[h(x)]^{s/r}\,dx. \tag{11}$$

If $x \in C_i$ or $\overline{C_i}$, using $|B(R)| = R^n|B(1)|$, we have

$$|B(|x|)|^\alpha \le \left\{ \begin{array}{cc} |B(1)|^\alpha 2^{in\alpha}, & \text{if } \alpha \ge 0, \\ |B(1)|^\alpha 2^{(i-1)n\alpha}, & \text{if } \alpha < 0, \end{array} \right\} \le |B(1)|^\alpha 2^{n|\alpha|}2^{in\alpha}, \tag{12}$$

and for $r > 0$,

$$|a+b+c|^r \le \left\{ \begin{array}{cc} |a|^r+|b|^r+|c|^r, & \text{if } 0<r\le 1, \\ 3^{r-1}(|a|^r+|b|^r+|c|^r), & \text{if } 1<r, \end{array} \right\} \le 3^r(|a|^r+|b|^r+|c|^r). \quad (13)$$

The first inequality is obvious when $0 < r \le 1$, from the property of convex function [14] when $r > 1$ since $g(x) = x^r$ is convex function. Noticing that $|b(x) - b(y)| \le |b(x) - b_{B_i}| + |b(y) - b_{\overline{B_j}}| + |b_{B_i} - b_{\overline{B_j}}|$, using (13) and (12), we have

$$\begin{aligned} h(x) &\le 2^{n|\alpha|}2^{n|\alpha-1|}3^r\frac{1}{2^{in\alpha}|B(1)|}\sum_{j=-\infty}^{i}2^{jn(\alpha-1)}\int_{\overline{C_j}}|b(x)-b_{B_i}|^r|f(y)|^r dy \\ &\quad +2^{n|\alpha|}2^{n|\alpha-1|}3^r\frac{1}{2^{in\alpha}|B(1)|}\sum_{j=-\infty}^{i}2^{jn(\alpha-1)}\int_{\overline{C_j}}|b(y)-b_{\overline{B_j}}|^r|f(y)|^r dy \\ &\quad +2^{n|\alpha|}2^{n|\alpha-1|}3^r\frac{1}{2^{in\alpha}|B(1)|}\sum_{j=-\infty}^{i}2^{jn(\alpha-1)}\int_{\overline{C_j}}|b_{B_i}-b_{\overline{B_j}}|^r|f(y)|^r dy \\ &= 2^{n|\alpha|}2^{n|\alpha-1|}3^r(I_1+I_2+I_3). \end{aligned}$$

By (10), we see that

$$\overline{C_j} \subset \overline{B_i} = B(|x|) \text{ for } j \le i \text{ and } x \in C_i; \text{ and } \frac{1}{2^{in}|B(1)|} \le \frac{1}{|\overline{B_i}|}.$$

Let $x \in C_i$, then we have

$$\begin{aligned} I_1 &\le |b(x)-b_{B_i}|^r\frac{1}{2^{in}|B(1)|}\int_{\overline{B_i}}|f(y)|^r dy\sum_{j=-\infty}^{i}2^{-(i-j)n(\alpha-1)} \\ &\le \frac{1}{1-2^{-n(\alpha-1)}}|b(x)-b_{B_i}|^r\frac{1}{|\overline{B_i}|}\int_{\overline{B_i}}|f(y)|^r dy, \end{aligned}$$

noticing that $\alpha - 1 > 0$. For I_2, since f is local bounded, by Holder's inequality and Lebesque's control convergence theorem, we have

$$\begin{aligned} \int_{\overline{C_j}}|b(y)-b_{\overline{B_j}}|^r|f(y)|^r dy &\le \left(\int_{\overline{C_j}}|b(y)-b_{\overline{B_j}}|^{rl'}dy\right)^{1/l'}\left(\int_{\overline{C_j}}|f(y)|^{rl}dy\right)^{1/l} \\ &\le |\overline{B_j}|^{1/l'}\|b\|_{CMO}^r\left(\int_{\overline{C_j}}|f(y)|^{rl}dy\right)^{1/l} \\ &\to \|b\|_{CMO}^r\int_{\overline{C_j}}|f(y)|^r dy, \quad (\text{when } l\to 1). \end{aligned} \quad (14)$$

Thus, as I_1,

$$I_2 \le \|b\|_{CMO}^r\frac{1}{1-2^{-n(\alpha-1)}}\frac{1}{|\overline{B_i}|}\int_{\overline{B_i}}|f(y)|^r dy.$$

For I_3, by (10),

$$\begin{aligned}
|b_{B_i} - b_{\overline{B_j}}| &\leq \frac{1}{|\overline{B_j}|}\int_{\overline{B_j}} |b(x) - b_{B_i}|dx \\
&\leq \begin{cases} \frac{1}{|B_j|}\int_{B_j} |b(x) - b_{B_i}|dx, & \text{if } j \leq i-1, \\ \frac{1}{|B_{j-1}|}\int_{B_j} |b(x) - b_{B_i}|dx, & \text{if } j = i, \end{cases} \\
&\leq \frac{2^n}{|B_j|}\int_{B_j} |b(x) - b_{B_i}|dx \\
&\leq \frac{2^n}{|B_j|}\int_{B_j} |b(x) - b_{B_j}|dx + 2^n \sum_{h=j}^{i-1} |b_{B_h} - b_{B_{h+1}}| \\
&\leq 2^{2n}\|b\|_{CMO}(i-j),
\end{aligned}$$

thus

$$\begin{aligned}
I_3 &\leq \|b\|_{CMO}^r \frac{1}{2^{in}|B(1)|}\int_{\overline{B_i}} |f(y)|^r dy \sum_{j=-\infty}^{i} 2^{-(i-j)n(\alpha-1)}(i-j)^r \\
&\leq c\|b\|_{CMO}^r \frac{1}{|\overline{B_i}|}\int_{\overline{B_i}} |f(y)|^r dy,
\end{aligned}$$

where $c = 2^{2nr}\sum\limits_{j=-\infty}^{i} 2^{-(i-j)n(\alpha-1)}(i-j)^r$. When $x \in C_i$, let

$$g(x) = \left(\frac{1}{|\overline{B_i}|}\int_{\overline{B_i}} |f(y)|^r dy\right)^{1/r} = \left(\frac{1}{|B(|x|)|}\int_{B(|x|)} |f(y)|^r dy\right)^{1/r},$$

combining to the estimates of I_1, I_2, I_3 above, and noticing that $(|b(x) - b_{B_i}|^r + \|b\|_{CMO}^r)^{s/r} \leq 2^{s/r}(|b(x) - b_{B_i}|^s + \|b\|_{CMO}^s)$ for $s > 0, r > 0$, we see that

$$\int_{C_i} |B(|x|)|^{\gamma-1}\,[h(x)]^{s/r}\,dx \leq c_0 \int_{C_i} |B(|x|)|^{\gamma-1}\,[|b(x) - b_{B_i}|^s + \|b\|_{CMO}^s]\,g^s(x)dx, \quad (15)$$

where $c_0 = [2^{n|\alpha|}2^{n|\alpha-1|}3^r \times 4 \times 2^{2nr}\sum\limits_{j=-\infty}^{i} 2^{-(i-j)n(\alpha-1)}(i-j)^r]^{s/r}$. While, since f local bound implies g local bound, as (14),

$$\int_{C_i} |B(|x|)|^{\gamma-1}|b(x) - b_{B_i}|^s g^s(x)dx \leq \|b\|_{CMO}^s \int_{C_i} |B(|x|)|^{\gamma-1} g^s(x)dx. \quad (16)$$

Thus, combining to (11), (15) and (16), we obtain

$$\begin{aligned}
M_s((M_{r,b}(f,\alpha),\gamma)(R) &\leq c_0^{1/s}\|b\|_{CMO}\left(\frac{1}{|B(R)|^\gamma}\sum_{i=-\infty}^{N}\int_{C_i} |B(|x|)|^{\gamma-1} g^s(x)dx\right)^{1/s} \\
&= c_0^{1/s}\|b\|_{CMO} M_s((M_r(f,1),\gamma)(R).
\end{aligned}$$

Using (1), we obtain (5).

The proof of (6). Replace (9) and (10) by

$$B_i = \begin{cases} B(2^i), & \text{if } i \geq N, \\ B(R), & \text{if } i = N-1, \end{cases}, C_i = B_i \setminus B_{i-1}, i = N, N+1, ..., \infty;$$

and if $x \in C_i$,

$$\overline{B_j} = \begin{cases} B_j, & \text{if } j \geq i, \\ B(|x|), & \text{if } j = i-1, \end{cases}, \overline{C_j} = \overline{B_j} \setminus \overline{B_{j-1}}, j = i, i+1, ..., \infty.$$

Then,

$$h^*(x) = \left[(M^*_{r,b}(f,\alpha)(|x|)\right]^r = \frac{1}{|B(|x|)|^\alpha} \sum_{j=i}^{\infty} \int_{\overline{C_j}} |B(|y|)|^{\alpha-1} (|b(x)-b(y)||f(y)|)^r dy,$$

and

$$\left[M^*_s((M^*_{r,b}(f,\alpha),\gamma)(R)\right]^s = \frac{1}{|B(R)|^\gamma} \sum_{i=N}^{\infty} \int_{C_i} |B(|x|)|^{\gamma-1} \left[h^*(x)\right]^{s/r} dx.$$

The rest of the proof of (6) is exactly similar to that of (5).

Thus, we finish the proof of Theorem 3.

The proofs of Theorem 4 by using (5)(6) are exactly similar to that of Theorem 2 by using (1)(2).

References

[1] A. Beurling, Construction and analysis of some convolution algebras, *Ann. Inst. Fourier (Grenoble)* **14**, (1964), 1-32.

[2] S. Bloom, A commutator theorem and weighted BMO, *Trans. Amer. Math. Soc.* **292**(1985), 103-122

[3] A.P. Calderon, Commutators ofsingular integral operators, Proc. Natl. Acad. Sci. USA 53 (1965) 1092-1099.

[4] R.Coifman, P. L. Lions, Y. Meyer and S. Semmes, Compensated compactness and Hardy spaces, *J. Math. Pures. Appl.* **72**, (1993),247-286.

[5] R. R. Coifman, R. Rochberg and G. Weiss, Factorization theorems for Hardy spaces in several variables variables, $Ann. of Math.$ **103**(1976), 611-635.

[6] Y. Z. Chen and K. S. Lau, Some new classes of Hardy spaces, *J.Functional Anal.* **84**(1989), 255-278.

[7] A. Cizmesija, J. Pecaric and I. Peric, Mixed means and inequalities of Hardy and Levin-Cochran-Lee type for multidimensional balls, *Proc. Amer. Math. Soc.* **128**, No.9 (2000), 2543-2552.

[8] A. Cizmesija and J. Pecaric, Mixed means and Hardy's inequality, *Math. Inequal. Appl.* **1**, No.4 (1998), 497-506.

[9] M. Christ, and L. Grafakos, Best constants for two nonconvolution inequalities, *Proc. Amer. Math. Soc.* **123**, No.6 (1995), 1687-1693.

[10] P. Drabek, H. P. Heinig and A. Kufner, Higher dimensional Hardy's inequality, *Int. Ser. Num. Math.* **123**(1997), 3-16.

[11] Z.Fu, Z.Liu, S.Lu, H.Wang, Characterization of the commutators of n-dim fractional order Hardy operator; Science in China Series A**37**, No.6, (2007), 651-659.

[12] J. Garcia-Cuerva, Hardy spaces and Beurling algebras, *J.LondonMath.Soc.* **39**(1989), 499-513.

[13] J. Garcia-Cuerva and M. J. L. Herrero, A theory of Hardy spaces associated to the Herz spaces, *Proc.LondonMath.Soc.* **69**, No.3(1994), 605-628.

[14] G. H. Hardy, J. E. Littlewood, and G. Polya , Inequalities, Cambridge Univ.Press, Cambridge, UK, 1959.

[15] S. Janson, Mean oscillation and commutators of singular integral operators, *Ark. Math.* **16**, (1978), 263-270.

[16] Y. Komori, Notes on commutators of Hardy operators [M];Intern J Pure Appl Math, **1**, (2003), 329-334.

[17] J. D. Lakey, Constructive decomposition of functions of finite central mean oscillation, *Proc. Amer. Math. Soc.* **127**, No.8 (1999), 2375-2384.

[18] Long Shunchao and Wang Jian, Commutators of Hardy operators, *J. Math. Anal. Appl.* **274**, No.2 (2002), 626-644.

[19] C. Perez, Endpoint estimates for commutators of singular integral operators, *J.Functional Anal.* **128**(1995), 163-195.

[20] M. Paluszynski, Characterization of the Besov spaces via commutators operator of Coifman, Rochberg and Weiss, *Indiana Univ. Math. J.* **44**, (1995), 1-17.

[21] Carlos Segovia and Jose L. Torrea, Higher order commutators for vector-valued Calderon-Zygmund operators, *Trans. Amer. Math. Soc.* **336**(1993),537-556.

[22] N. Wiener, Generalized harmonic analysis, *Acta Math* . **55**, (1930), 117-258.

[23] A. Youssfi, Regularity properties of commutators and BMO-Triebel-Lizorkin spaces, *Ann. Inst. Fourier, Grenoble.* , No.3 (1995), 795-807.

Long Shunchao, Jian Wang,
Mathematics Department,
Xiangtan University,
Xiangtan , 411105, China,
E-mail address: sclong@xtu.edu.cn